



AGRONOMIC SPOTLIGHT



UNDERSTANDING SEED TREATMENTS FOR SWEET CORN

- » Seed treatments protect sweet corn seed and seedlings from soilborne and seedborne pathogens and insect pests.
- » Seed treatments are especially important for supersweet (SH2) hybrids.
- » A range of fungicides is needed to control a variety of sweet corn pathogens.

Recommendations for treating sweet corn seed date back as far as 1926,¹ and seed treatments are still a very effective means of protecting seeds and seedlings from soilborne and seedborne pathogens and insect pests. They also help increase rapid emergence, optimize stand establishment, and promote strong root growth. Seed treatments are especially beneficial for supersweet (SH2) hybrids because these hybrids tend to germinate more slowly and grow less vigorously than those of standard-sugary (su) hybrids.²

SEED AND SEEDLING DISEASES

Soilborne and seedborne pathogens can infect sweet corn seeds and seedlings, resulting in reduced stands and seedling vigor. Seedborne pathogens include the fungi *Fusarium moniliforme*, *Penicillium oxalicum*, and species of *Rhizopus*.² Soilborne pathogens include *Pythium ultimum*, *Rhizoctonia solani*, and several species of *Fusarium*. Infection by these pathogens often results in pre- or post-emergence damping off. Damping-off is favored by cool, wet conditions that slow germination and seedling growth.

STANDARD SWEET CORN SEED TREATMENTS

The International Sweet Corn Development Association (ISCDA) Seed Treatment Committee was formed in the early 1990s to help the sweet corn industry identify the best performing seed treatments for sweet corn through coordinated trials carried out at locations across the United States. These trials have shown that effective seed treatments should include one or more broad-spectrum fungicides, systemic fungicides with activity against seedborne *Penicillium* and *Fusarium*, and a fungicide with activity against Oomycete pathogens such as *Pythium* and *Phytophthora*. In locations where insect pests are present, the inclusion of an insecticide is also beneficial.³

THE NEED FOR FUNGICIDES

In a 2012 ISCDA coordinated study, 19 seed treatment combinations, were evaluated at 16 different locations.² The study included two product combinations representing industry standard treatments; **standard treatment 1** (Dividend Extreme[®], Apron[®] XL, Maxim[®] 4FS, Vitavax[®] -34)

TABLE 1. Effects of standard seed treatments and a non-treated control treatment for stand establishment and seedling vigor of sweet corn.²

Seed Treatment	Group 1 trials (cool, wet at planting)		Group 2 trials (normal conditions)	
	Stand Count (%)	Vigor*	Stand Count (%)	Vigor
Non-treated control	34.2 b**	2.53 b	57.9 c	2.95 b
Standard treatment 1: Dividend Extreme [®] Apron [®] XL, Maxim [®] 4FS, Vitavax [®] -34	71.1 a	3.64 a	69.0 b	3.53 a
Standard treatment 2: captan, thiram, Dividend Extreme [®] , Apron [®] XL, Vitavax [®] -34	73.7 a	3.55 a	74.1 a	3.77 a


* Plant vigor was rated on a scale from 1 to 5 at the 5-6 leaf stage.

** Values followed by the same letter within a column are not significantly different.

The data presented here is a subset of a larger set of results reported in the referenced article.²

and **standard treatment 2** (captan, thiram, Dividend Extreme[®], Apron[®] XL, Vitavax[®] -34). In this study, stand counts and seedling vigor were significantly higher in plots planted to seed treated with either of the two standard combinations as compared to plots planted with non-treated seed (Figure 1). In locations with cool, wet soil conditions, the two standard combinations performed equally well, but in locations with warmer, drier conditions, the combination containing captan and thiram resulted in higher stand counts and levels of seedling vigor (Table 1).

Damping-off caused by *Pythium* can be a significant problem



with sweet corn. As a result, most sweet corn seed is treated with fungicides that contain metalaxyl or mefenoxam. The widespread

Figure 1. Seed treatments can improve both stand establishment and seedling vigor. Stand establishment in an ISCDA trial; (a) no treatment, (b) standard treatment 1, (c) standard treatment 2.² Carrie Wohleb, Washington State University.

(Continued on page 2)





UNDERSTANDING SEED TREATMENTS FOR SWEET CORN

(Continued from page 1)

use of this fungicide class increases the likelihood of the development of *Pythium* strains that are resistant to these chemicals. In recent years, metalaxyl-resistant isolates of *Pythium ultimum* have been found in the Pacific Northwest. This emphasizes the need for the availability of products with different modes of action against oomycete pathogens. For the past several years the ISCTA trials have evaluated the efficacy of fungicides for their activity against organisms such as *Pythium* in order to identify alternate products that could be used instead of metalaxyl/mefenoxam. Ethaboxam (Intego™ Solo Fungicide) is a newer fungicide that was found to be effective against *Pythium* damping-off in recent ISCTA seed treatment trials.²

THE NEED FOR INSECTICIDES

Soil insects, such as wireworm, seedcorn maggot, white grub, and grape colaspis, can feed on seed kernels and destroy germinating seeds. These insects can be especially problematic when sweet corn is planted in weedy fields previously fallow or planted to turf. Under these conditions, insecticide seed treatments can be beneficial.

The insecticides used most commonly for seed treatment in sweet corn belong to the neonic (short for nitroguanidine neonicotinoid) class of insecticides. These are systemic insecticides that are effective against a wide range of insect pests, but they are less toxic to birds and mammals than organophosphate and carbamate insecticides. Neonic products used in seed treatments include clothianidin (Poncho® 600), imidacloprid (Gaucho® 600), and thiamethoxam (Cruiser® 5FS).

If populations of soil insect pests are at or above threshold levels, soil applied insecticides (e.g. phorate, terbufos, chlorpyrifos, ethoprop) may be needed in addition to the insecticide seed treatment to provide adequate control.⁴

In addition to controlling insect pests, seed treatment insecticides can also reduce levels of insect-vectored diseases, such as Stewart's wilt, a bacterial disease vectored by corn flea beetles.⁵

SEMINIS SEED TREATMENT OPTIONS

Seminis currently offers several sweet corn seed treatments containing various combinations of products used to control fungal pathogens and insect pests. Table 2 lists the fungicide and insecticide products and describes what they are intended to control.



TABLE 2. FUNGICIDE AND INSECTICIDE PRODUCTS FOUND IN SEMINIS® SWEET CORN SEED TREATMENTS AND THE DISEASES AND PESTS CONTROLLED.

Product (active ingredient)	Protects against	Treatments *				
		Conventional			Performance Series®	
		S	P	AP	PSS	PSP
Apron XL® (mefenoxam)	damping-off caused by <i>Pythium</i> and <i>Phytophthora</i>	✓	✓	✓	✓	✓
Dividend Extreme® (mefenoxam + difenoconazole)	damping-off caused by <i>Pythium</i> , <i>Phytophthora</i> , and seedborne <i>Penicillium</i>	✓	✓	✓	✓	✓
42-S Thiram	a broad-spectrum fungicide; protects from several soilborne fungi	✓	✓	✓	✓	✓
Vitavax®-34 (carboxin)	seedborne head smut, seed decay, and damping-off fungi, including <i>Rhizoctonia solani</i>	✓	✓	✓	✓	✓
Maxim® 4FS (fludioxonil)	seedborne and soilborne fungi that cause seed decay and damping-off, including <i>Rhizoctonia solani</i> and several species of	✓	✓	✓	✓	✓
Dynasty® (azoxystrobin)	seedborne and soilborne fungi that cause decay, damping-off and seedling blight, including species of <i>Rhizoctonia</i> , <i>Penicillium</i> , and <i>Pythium</i> , and some protection from seedborne head smut		✓		✓	
Poncho®600 (clothianidin)	chinch bug, corn flea beetle, corn leaf aphid, cutworm, grape colaspis, seedcorn maggot, southern corn leaf beetle, southern corn rootworm, southern green stinkbug, sugarcane beetle, white grub, thrips, and wireworms		✓		✓	
Cruiser®5FS (thiamethoxam)	wireworm, seed corn maggot, leaf beetle, white grub, black cutworm, thrips, green stinkbug, seed corn beetle, corn leaf aphid, sugarcane beetle, corn rootworm, and billbug			✓		✓

* Seminis seed treatment options: Conventional refers to treatments available for conventional sweet corn hybrids; Performance Series® refers to treatments available on Performance® Series sweet corn. S=Standard (conventional), P=Premium (conventional), AP=Alternate Premium (conventional), PSS=Standard (Performance Series®), PSP=Premium (Performance Series®)

Sources:

- ¹ Reddy, C.S., Holbert, J.R., and Erwin, A.T. 1926. Seed treatments for sweet-corn diseases. Journal of Agricultural Research 33:769-779.
- ² Wohleb, C.H. 2014. Evaluation of fungicide seed treatments for sweet corn. Journal of the NACAA Volume 7.
- ³ Wohleb, C.H. 2013. The ISCTA multi-location seed treatment trials for sweet corn. Pacific Northwest Vegetable Conference November 13, 2013.
- ⁴ Mossler, M.A. 2014. Crop profile for sweet corn in Florida. University of Florida IFAS Extension. CIR1233.
- ⁵ Pataky, J.K., Michener, P.M., Freeman, N.D., Weinzierl, R.A., and Teyker, R.H. 2000. Control of Stewart's wilt in sweet corn with seed treatment insecticides. Plant Dis. 84:1104-1108.

IMPORTANT: Produce Marketing and Stewardship Requirements: This product has been approved for import into key export markets with functioning regulatory systems. Any crop or material produced from this product can only be exported to, or used, processed or sold in countries where all necessary regulatory approvals have been granted. It is a violation of national and international law to move material containing biotech traits across boundaries into nations where import is not permitted. It is the grower's responsibility to talk to their produce handler or purchaser to confirm their buying position for this produce so that the marketing requirements can be met.

Herbicide Information for Performance Series® sweet corn: Roundup PowerMAX®, Roundup PowerMAX® II and Roundup WeatherMAX® herbicides are approved for use on Performance Series® sweet corn (containing the Roundup Ready® trait) in all U.S. states, the District of Columbia and Puerto Rico. If the directions for use on sweet corn with Roundup Ready® 2 Technology (which includes Performance Series® sweet corn) are not listed in the product label that is attached to the product you purchased, contact your Monsanto Company representative.

Performance Series® sweet corn Insect Resistance Management (IRM) – Post-Harvest Requirements: Crop destruction must occur no later than 30 days following harvest, but preferably within 14 days. The allowed crop destruction methods are: rotary mowing, discing, or plowing down. Crop destruction methods should destroy any surviving resistant insects.

B.t. products may not yet be registered in all states. Check with your Monsanto representative for the registration status in your state. All information concerning Performance Series® sweet corn hybrids given orally or in writing by Monsanto or its employees or agents, including the information in this communication, is given in good faith, but is not to be taken as a representation or warranty by Monsanto as to the performance or suitability of Performance Series® sweet corn hybrids, which may depend on local climatic conditions and other factors. Monsanto assumes no liability for any such information. This information shall not form part of any contract with Monsanto unless otherwise specified in writing. 161028150112 012017DME

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Roundup Ready® 2 Technology contains genes that confer tolerance to glyphosate, an active ingredient in Roundup® brand agricultural herbicides. Agricultural herbicides containing glyphosate will kill crops that are not tolerant to glyphosate. Performance Series®, Roundup PowerMAX®, Roundup Ready 2 Technology and Design®, Roundup Technology® and Roundup WeatherMAX® are trademarks of Monsanto Technology LLC. Seminis® is a registered trademark of Seminis Vegetable Seeds, Inc. All other trademarks are the property of their respective owner. ©2017 Seminis Vegetable Seeds, Inc.

